203

- 23. (2x Amended) A method for potentiating an antigen-specific immune response in a subject comprising:
- a. pretreating an area of skin of said subject, whereby said pretreating disrupts at least the skin's stratum corneum but does not penetrate the skin's dermis;
- b. applying an adjuvant formulation to said pretreated area, wherein said adjuvant formulation comprises:
- 1) at least one adjuvant present in an amount effective to promote said immune response and
- 2) a pharmaceutically acceptable carrier, wherein said pretreating enhances skin penetration by said adjuvant formulation; and
- c. administering to said subject a separate antigen formulation comprising at least one antigen sufficient to induce an antigen-specific immune response against a pathogen; such that said at least one antigen is processed by an antigen presenting cell and then presented to a lymphocyte to induce said immune response, and said at least one adjuvant potentiates said immune response.
- 28. (2x Amended) A method for potentiating an antigen-specific immune response in a subject comprising:
- a. pretreating an area of the skin of said subject, whereby said pretreating disrupts at least the skin's stratum corneum but does not penetrate the skin's dermis;
- b. applying an antigen formulation to said pretreated area, wherein said antigen formulation comprises:
- 1) at least one antigen sufficient to induce an antigen-specific immune response against a pathogen and
- 2) a pharmaceutically acceptable carrier, wherein said pretreating enhances skin penetration by said antigen formulation; and
- c. administering to said subject a separate adjuvant formulation comprising at least one adjuvant in an amount effective to promote said immune response;

GLENN et al. - Appln. No. 09/257,188

DRAFT

such that said at least one antigen is processed by an antigen presenting cell and then presented to a lymphocyte to induce said immune response, and said at least one adjuvant potentiates said immune response.

Kindly add the following new claims.

- 92. (New) The method of claim 23, wherein said separate antigen formulation is administered at said pretreated area of skin.
- 93. (New) The method of claim 23, wherein said separate antigen formulation is administered at a site different from said pretreated area of skin.
- 94. (New) The method of claim 28, wherein said separate adjuvant formulation is administered at said pretreated area of skin.
- 95. (New) The method of claim 28, wherein said separate adjuvant formulation is administered at a site different from said pretreated area of skin.